

form “OR-AI”

Lansford

Injection Laryngoplasty

👉 Preferences, Pick Sheet, and

“Anything Special?”

Note: This is a different procedure from External Implant Medialization (Isshiki) Thyroplasty, which uses form “OR-AJ”

Color code:

Surgeon
Anesthesia
Circulating Nurse
Scrub Tech
Surgical Assistant (Not necessary for this case)



QR for online color version. Select Form “OR-AI”

Pre-op

Surgeon signs chart, and reviews his own consent form with patient and a witness **prior to** administration of an anxiolytic or other medication rendering a patient incompetent to sign. Minors may receive an anxiolytic prior to the surgeon’s consent since they are not signing.

Instruments & Supplies

*** The substance to be injected may need to be ordered prior to the day of surgery ***
Ask Dr. Lansford which material is needed. Common injection substances include:
Renu
ProLaryng
Restylane (Q-Med, Uppsala, Sweden)
Surgiderm 24 XP (Allergan)
Hylan B gel (Hylaform, Biomatrix, Ridgefield, NJ, USA)
At least 1 mL is needed.

Laryngoscopy Tray
25 ga laryngeal needle
or, alternatively 25 ga butterfly needle
Suspension apparatus
0 degree and 30 degree rigid telescope (sinus scopes)
Camera and light carriers for scopes
Ortho Pack
Suction tubing
Mayo cover x 2
1/2” x 1/2” cottonoids

Defog pad
Teeth guard (maybe use one from anesthesia)
Size 8 Sensicare gloves
Lidocaine (1% or 2%) with epinephrine (1:200k up to 1:100k)
Head ring/donut (gel preferred)
Shoulder roll (gel preferred)
Small ETT (5.5 or smaller for adults)

Equipment

Video tower
Chair with adjustable height

Patient in Room to Induction

Pt. Supine, head at the tippy-top of the bed
Turn bed 90 degrees, reverse Trendelenburg
Paralysis is usually needed
SCDs

From Induction to Start

Head wrap - two towels and a towel clip
Towels and split sheet, but anesthesia circuit exposed

Mid Operation

Music and background discussion quiet, please.
Possible specimens for pathology - usually in formalin
Mild hypotension, such as SBP ~105 mmHg, if tolerated.

Closing

Let the surgeon know if you want lidocaine on the vocal cords
Communicate with the surgeon regarding any airway concerns

Emergence to Patient Exit

Try to minimize patient coughing

