

form “OR-Y”

Lansford Submandibular Gland Excision



QR for online color
version. Select
Form “OR-Y”

👉 Preferences, Pick Sheet, and “Anything Special?”

Color code:

Surgeon
Anesthesia
Circulating Nurse
Scrub Tech
Surgical Assistant

Pre-op

Surgeon signs chart, and reviews his own consent form with patient and a witness **prior to** administration of an anxiolytic or other medication rendering a patient incompetent to sign. Minors may receive an anxiolytic prior to the surgeon’s consent since they are not signing.

Unsterile Prep Tray

Shave neck if needed.
Regular tip (not fine) skin marking pen and ruler
1% lidocaine with 1:100,000 epinephrine in a 10 mL syringe and a 27 ga, 1.5” needle
Wet Betadine prep tray scrub and paint (or chloraprep if allergic)
4x4” gauze (or similar)

Instruments & Supplies

Neck dissection tray
Bipolar cautery - bayonet forceps, guarded
Bovie pencil and guarded flat blade tip
Large “U” split drape
Stapler
Size 8 Sensicare gloves for CDL
XL , XLNG surgical gown for CDL
#15 scalpel blades
Tenotomy scissors

Debakey x 2
Double prong skin hooks, wide x 2
Weitlander, hinged, medium x 2
Andrews metal Yankauer suction tip
Army-Navy retractors x 2
Right angle hemostat clamps x 2
Kitners (1 pack)
Saline for irrigation
10 mL syringe
2-0 silk on RB-1 x 2
3-0 Vicryl SH x 2
4-0 Monocryl on PS-2 x 1
5-0 Prolene, blue x 1
JP drain reservoir (have in room unopened)
10 Fr slotted Blake channel drain on a trocar(have in room unopened)
Wound dressings:
 Bacitracin
 Xeroform strip
 Telfa
 Tegaderms - 2 small, 2 medium

Equipment

Headlight
Electrocautery (Bovie)
holder roll (gel preferred)
Head ring/donut (gel preferred)
Pump chair without arms (SurgiStool II)
Stryker bed with adjustable angle head board at the top of the bed
Bair Hugger from nipples down
Fluid warmer on bed
SCDs
Pillow for under knees
OR Neptune
Extension ("pediatric") airway circuit tubing

Patient in Room to Induction

Will turn OR table 180 degrees
Adjustable head board at top of bed
Position patient at tippy top of bed
ETT taped to opposite side
Pre-op antibiotic within 60 minutes before incision
No paralysis during case
Decision for foley per anesthesia service; typical OR time 60-90 min.

From Induction to Start

Arm on operative side tucked with foam
10 mL of 1-2% lido with epi 1:100,000 (or similar) with 27 ga needle.
Doctor will inject local prior to prep.

Bair Hugger for lower body below xiphoid
Betadine paint - From nose to below the nipple line, right face, neck, chest but crossing
midline. Try not to erase markings.
Bovie set to 15 cut/15 coag

Mid Operation

Music and background discussion quiet, please.
Specimen for pathology- usually routine pathology in formalin, but ASK
Mild hypotension, such as SBP ~105 mmHg, if tolerated.
Keep a squared Ray-tec on the field at all times

Closing

Bacitracin, Xeroform strip, Telfa, Tegaderm for the incision.

Emergence to Patient Exit

Thank you for reading this!